

Master Equations for Extended Lagrangian BRST Symmetries

POUL H. DAMGAARD and FRANK DE JONGHE*

CERN – Geneva

February 1, 2008

Abstract

Starting from the requirement that a Lagrangian field theory be invariant under both Schwinger-Dyson BRST and Schwinger-Dyson anti-BRST symmetry, we derive the BRST–anti-BRST analogue of the Batalin-Vilkovisky formalism. This is done through standard Lagrangian gauge fixing respecting the extended BRST symmetry. The solutions of the resulting Master Equation and the gauge-fixing procedure for the quantum action can be brought into forms that coincide with those obtained earlier on algebraic grounds by Batalin, Lavrov and Tyutin.

hep-th@xxx/9302050

KUL-TF-93/04

CERN-TH-6800/93

February 1993

*Aspirant of the N.F.W.O., Belgium. On leave from the Instituut voor Theoretische Fysica, University of Leuven, Belgium.

Schwinger-Dyson equations are believed to provide a complete description of any quantum field theory, once appropriate boundary conditions are imposed. To implement the content of these Schwinger-Dyson equations as a dynamical principle in Lagrangian quantization, one needs the symmetry operator that enforces the equations. This “Schwinger-Dyson BRST operator”, in the following denoted by δ , was first found in ref. [1]. (For some generalizations, see ref. [2]).

Recently, it was discovered [3] that demanding a Lagrangian field theory to be invariant under this Schwinger-Dyson BRST symmetry in addition to usual BRST symmetries of internal gauge invariances leads directly to the Batalin-Vilkovisky Lagrangian formalism [4]. The derivation is done most simply in terms of a certain Lagrangian collective field formalism, described in detail in ref. [2], but can also be based directly on the Schwinger-Dyson BRST symmetry itself.

One interesting aspect of these observations is that the Batalin-Vilkovisky formalism, normally based on a one-to-one doubling of field variables¹ (to each fundamental “field” ϕ^A , including usual ghosts and antighosts whenever needed, one associates an “antifield” ϕ_A^*), actually corresponds to an intermediate step of the Lagrangian quantization procedure, in which a certain set of ghost fields c^A already have been integrated out. These new ghosts c^A play a fundamental rôle in the Schwinger-Dyson BRST transformations [1]. When the functional measures for the fundamental fields are flat, the Schwinger-Dyson BRST algebra reads (our conventions follow those of ref. [3]):

$$\begin{aligned}\delta\phi^A(x) &= c^A(x) \\ \delta c^A(x) &= 0 \\ \delta\phi_A^*(x) &= -\frac{\delta^l S}{\delta\phi^A(x)} ,\end{aligned}\tag{1}$$

with S generically indicating the full action. The Ward Identities following from this symmetry, when implemented in the Lagrangian path integral, are precisely the most general Schwinger-Dyson equations. The way to include internal gauge symmetries of the starting point, the given classical action S_0 , is described in ref. [3]. After integrating out the new ghosts c^A , the method is completely equivalent to the Batalin-Vilkovisky scheme [4]. The antifields ϕ_A^* are simply the usual antighosts of the ghosts c^A that enforce Schwinger-Dyson equations through shifts of the fundamental fields.

Algebraically, the Schwinger-Dyson BRST transformations provide a symmetry of the full quantum partition function whenever the quantum action S_{ext} satisfies a Master Equation of the form

$$\frac{1}{2}(S_{ext}, S_{ext}) = -\frac{\delta^r S_{ext}}{\delta\phi^A} c^A + i\hbar\Delta S_{ext} ,\tag{2}$$

with the antibracket $(\cdot, \cdot)$ and the operator Δ coinciding with those of Batalin and Vilkovisky [4]. It is important to realize that the above Master Equation must be imposed from the beginning, independently of any possible gauge fixings. For theories of flat measures (and we shall consider only those cases here), eq. (2) is a necessary condition

¹For some excellent recent reviews of the Batalin-Vilkovisky formalism, see ref. [5].

that the final quantum field theory gives rise to the correct Schwinger-Dyson equations. Gauge-independence of the generating functional upon any subsequent gauge fixings follows straightforwardly as a by-product [3].

A sufficient condition to guarantee that the resulting Ward Identities of S_{ext} contain all Schwinger-Dyson equations for gauge-invariant Green functions of the classical fields is that the solution S_{ext} of eq. (2) can be written

$$S_{ext}[\phi, \phi^*, c] = S^{(BV)}[\phi, \phi^*] - \phi_A^* c^A, \quad (3)$$

with $S^{(BV)}$ then required to satisfy the quantum Master Equation of Batalin and Vilkovisky:

$$\frac{1}{2}(S^{(BV)}, S^{(BV)}) = i\hbar \Delta S^{(BV)}. \quad (4)$$

Upon integrating out c^A , this automatically sets $\phi_A^* = 0$ before gauge fixing. This also, together with the Batalin-Vilkovisky boundary conditions [4] imposed on $S^{(BV)}$, guarantees that the final gauge fixing is achieved by a certain canonical transformation (within the antibracket).

The resulting Lagrangian formalism corresponds directly to (but provides a much more powerful machinery than) standard BRST Lagrangian quantization. In particular, the action S_{ext} itself can in a certain specific sense [4] be viewed as the Lagrangian BRST generator.

As formulated above, this seems to leave out the anti-BRST symmetries that can always be constructed for closed algebras, given the BRST symmetries [6]. However, it was shown already in ref. [1] that there also exists a corresponding *Schwinger-Dyson anti-BRST symmetry*. The question then naturally arises whether there also exists a corresponding Lagrangian anti-BRST generator. In particular, can one set up a corresponding quantization prescription that leads to a path integral invariant under *extended* BRST symmetry, *i.e.* simultaneous invariance under both BRST and anti-BRST symmetries? This question has been answered in the affirmative by Batalin, Lavrov and Tyutin [7] from a completely algebraic starting point.² For those who find the standard Batalin-Vilkovisky scheme (with its doubling of fields and subsequent conditions on how to remove the antifields) rather formidable, the extended BRST scheme of Batalin, Lavrov and Tyutin must appear even more impressive. It requires the introduction of now *three* antifields ϕ_{Aa}^* , $a = 1, 2$ and $\bar{\phi}^A$, all of which are finally removed through a certain gauge-fixing procedure that freezes their values [7].

If the ordinary Batalin-Vilkovisky Lagrangian scheme follows from imposing the Schwinger-Dyson BRST symmetry on the path integral, one would expect that the most natural starting point for deriving a path integral invariant under extended BRST symmetries would be to impose simultaneously invariance under both the Schwinger-Dyson BRST *and* Schwinger-Dyson anti-BRST symmetries. Because the algebra corresponding to the Schwinger-Dyson anti-BRST symmetry [1] can be formulated in terms of the same multiplet of fields needed for the corresponding BRST symmetry, one would a priori not expect new fields to appear. One purpose of the present letter is to clarify that issue.

²For some alternative formulations, see also ref. [8].

Starting from the same collective field technique employed in ref. [3], we shall *derive* the Master Equation that gives rise to an action of extended BRST symmetry in the case of an irreducible closed algebra.

We start by recapitulating the BRST and anti-BRST transformations for Schwinger-Dyson equations [1]. We will change notation and denote the BRST operator by δ_1 and its anti-BRST companion by δ_2 . As we wish to treat both symmetries on an equal footing, which means that the ghosts c^A are viewed as $Sp(2)$ partners of the antighosts ϕ_A^* , we choose a more symmetric notation in which c^A and ϕ_A^* are considered as the first and second component of a combined field ϕ_{Aa}^* , $a = 1, 2$.

In order to see the extended Schwinger-Dyson symmetries and their consequences for the action, it suffices to consider a theory of classical action $S[\phi]$ that is free of internal gauge invariances. Introducing collective fields $\bar{\phi}^A$ for the ϕ^A -fields³, we gauge the shift symmetry of the flat ϕ^A -measure through the transformation $\phi^A \rightarrow \phi^A - \bar{\phi}^A$, and integrate over these collective fields $\bar{\phi}^A$ as well. The fields ϕ^A are of arbitrary Grassmann parity ϵ_A , and we of course require $\epsilon(\bar{\phi}^A) = \epsilon_A$. There is no distinction between upper and lower indices on ϕ . The BRST (and anti-BRST) versions of the above gauge symmetry are the sought-for Schwinger-Dyson BRST (and anti-BRST) symmetries [1]:

$$\begin{aligned}\delta_a \phi_A &= \phi_{Aa}^* \\ \delta_a \bar{\phi}^A &= \phi_{Aa}^* \\ \delta_a \phi_{Ab}^* &= \varepsilon_{ab} B_A \\ \delta_a B_A &= 0 ,\end{aligned}\tag{5}$$

where ε_{ab} is the $Sp(2)$ invariant tensor ($\varepsilon_{12} = 1, \varepsilon_{ab} = -\varepsilon_{ba}$).

It follows trivially that these transformations are both nilpotent and satisfy the correct BRST–anti-BRST algebra

$$\delta_a \delta_b + \delta_b \delta_a = 0 .\tag{6}$$

Using the fact that $gh(\delta_a) = -(-1)^a$ we derive the following ghost number assignments:

$$\begin{aligned}gh(\phi_{Aa}^*) &= -(-1)^a + gh(\phi^A) \\ gh(\bar{\phi}^A) &= gh(\phi^A) \\ gh(B^A) &= gh(\phi^A) .\end{aligned}\tag{7}$$

Grassmann parities of the new fields are $\epsilon(\phi_{Aa}^*) = \epsilon_A + 1$ and $\epsilon(B_A) = \epsilon_A$.

Gauge-fixing the collective fields to zero in a manner manifestly invariant under the extended Schwinger-Dyson BRST can be achieved by adding a term

$$\frac{1}{4} \varepsilon^{ab} \delta_a \delta_b [\bar{\phi}_A (-1)^{\epsilon_B} M_{AB} \bar{\phi}_B] = (-1)^{\epsilon_B+1} \bar{\phi}_A M_{AB} B_B - \frac{1}{2} \varepsilon^{ab} \phi_{Aa}^* M_{AB} \phi_{Bb}^* .\tag{8}$$

Here M_{AB} is a constant invertible matrix of symmetry $M_{AB} = (-1)^{\epsilon_A \cdot \epsilon_B} M_{BA}$. All elements of M between Grassmann odd and even sectors vanish. M ensures that the term we add

³The choice of notation for the collective fields ($\bar{\phi}^A$) differs from that of ref. [3]. The reason for the present choice will become obvious below.

to the action has overall ghost number zero, and is even under Grassmann parity.⁴

Integrating out $\bar{\phi}_A$ and B_A yields the action

$$S_{ext} = S[\phi] - \frac{1}{2} \varepsilon^{ab} \phi_{Aa}^* M_{AB} \phi_{Bb}^* , \quad (9)$$

which is invariant under the more compact version of the extended Schwinger-Dyson BRST symmetry

$$\begin{aligned} \delta_a \phi_A &= \phi_{Aa}^* \\ \delta_a \phi_{Ab}^* &= \varepsilon_{ba} M_{BA}^{-1} \frac{\delta^l S_{ext}}{\delta \phi_B} \end{aligned} \quad (10)$$

obtained by substituting for B_A the equation of motion. Ward Identities of the form $0 = \langle \varepsilon^{ab} \delta_b \phi_{Aa}^* F[\phi] \rangle$ are the most general Schwinger-Dyson equations

$$\left\langle \frac{\delta^l F}{\delta \phi_A(x)} + \left(\frac{i}{\hbar} \right) \frac{\delta^l S}{\delta \phi_A(x)} F[\phi_A] \right\rangle = 0 \quad (11)$$

of the partition function

$$\mathcal{Z} = \int [d\phi^A][d\phi_{Aa}^*] \exp \left[\frac{i}{\hbar} S_{ext} \right] . \quad (12)$$

Turning the argument around, we can now write down algebraic equations the extended action S_{ext} must satisfy in order to be invariant under the extended Schwinger-Dyson BRST symmetry (10). Inserting the transformations (10) into the conditions $0 = \delta_a S_{ext}$ is seen to be equivalent to demanding that S_{ext} must satisfy the following Master Equations:

$$\varepsilon_{ba} \frac{\delta^r S_{ext}}{\delta \phi_{Aa}^*} M_{BA}^{-1} \frac{\delta^l S_{ext}}{\delta \phi_B} = \frac{\delta^r S_{ext}}{\delta \phi^A} \phi_{Ab}^* . \quad (13)$$

This is just the $Sp(2)$ invariant version of the classical part of the Master Equation (2). Because of the derivation in terms of collective fields, we know that the functional measure (formally) remains invariant.⁵ We are throughout assuming the existence of a suitable regularization scheme that respects the BRST symmetry. If one insists on regulators that break the symmetry, the problem must be cast in terms of the quantum Master Equation, to be discussed briefly below. (Some aspects of this issue in the usual Batalin-Vilkovisky BRST framework are discussed in ref. [9].)

Next, we turn to the question of internal gauge symmetries. In the case of just BRST symmetry, it was found that the inclusion of gauge symmetries did not fundamentally alter the formalism. In particular, the Master Equation remains the same. This is as one

⁴This matrix M is also required to gauge-fix in a BRST-anti-BRST preserving manner Grassmann-odd fields to zero. Its presence may be avoided by the introduction of more fields in the extended BRST multiplet, but for our purposes the present formulation suffices. M will drop out in the end.

⁵The action and the starting measure are trivially invariant, and the combination of action and measure must respect the symmetry even after having integrated out some of the fields. The measure therefore must remain invariant. This can of course also be checked explicitly.

would expect, because Schwinger-Dyson equations should be satisfied independently of particular choices of gauge fixing. As we have mentioned above, a solution of the kind (3) ensures that the resulting equations for the original *classical* fields coincide with those of the original action S . Introducing the Batalin-Vilkovisky boundary conditions [4] implies that the final gauge fixing procedure can be done through a canonical transformation alone. Different boundary conditions can be equally valid for the case of closed algebras, but then the gauge-fixing procedure may require the usual addition of a BRST-exact term [3].

Imposing only anti-BRST symmetry can be done in a completely similar manner, and we shall not elaborate on it.

However, insisting on having *extended* BRST–anti-BRST symmetry brings in a fundamentally new aspect. In order to set up the gauge-fixing of the extended action S_{ext} satisfying the Master Equation (13), one must find the equivalent of adding a gauge-fixing term that is simultaneously invariant under both the BRST and anti-BRST symmetries. Normally, this is done by adding a term of the form $\varepsilon^{ab}\delta_a\delta_b\chi(\phi)$, for a suitable bosonic function χ . But after having integrated out the auxiliary fields B_A , the operator $\varepsilon^{ab}\delta_a\delta_b$ is no longer nilpotent, even when acting on the subspace of fields ϕ_A . This means that there is no simple way of starting directly with the Master Equation (13), introducing suitable boundary conditions, and then finding an appropriate way of gauge fixing.⁶

This problem is instantly cured if we revert to the formulation in terms of collective fields $\bar{\phi}_A$. The operator $\varepsilon^{ab}\delta_a\delta_b$ is now by construction nilpotent, and we can use it to add BRST–anti-BRST exact terms to the action without changing the physical content. But of course this also means that we are restricted to situations where the collective field formalism can be applied. In the case of open gauge algebras a consistent formalism that allows us to merge the collective field BRST symmetry with that of the internal gauge algebra has not yet been found. Thus by remaining at the level where the extended BRST symmetry is still formulated with the help of the collective fields discussed above (and no more), we are at present prevented from constructing, for open gauge algebras, a gauge-fixed action invariant under the extended BRST symmetries. It is not yet clear whether this implies that extended BRST symmetries cannot be imposed on such theories. Introducing more fields may be one way of extending properly the collective field formalism to that case.

The starting point, then, is that for a closed irreducible gauge algebra, the existence of both BRST and anti-BRST transformations [6] leads us to consider *two* nilpotent transformations induced by the internal gauge symmetry. As is customary, ϕ_A denotes collectively all classical fields, ghosts and antighosts, and auxiliary fields. The BRST and anti-BRST transformations for the internal symmetries can then be written in the general for $\delta_a\phi_A = \mathcal{R}_{\mathcal{A}+}$. These transformations satisfy (6) and $\delta_a\mathcal{R}_{\mathcal{A}+} = \mathbf{1}$ for both values of a . To end the cycle, one must introduce suitable Nakanishi-Lautrup fields, which are now forced to transform non-trivially. These fields will be integrated over in the path integral as well, but their precise transformation properties need not concern us if we simply include these

⁶This problem does not arise if one insists only on preserving either BRST or anti-BRST invariance. A sufficient gauge-fixing term is then of the form $\delta_a\Psi(\phi)$ for a certain fermionic function Ψ , and the operator δ_a is nilpotent when acting only on the subspace of ϕ_A , – which is all that is required.

fields in the ϕ^A . The explicit form of $\mathcal{R}_{A+}$ in the case of an irreducible closed algebra can be found in ref. [6].

We can now straightforwardly combine these extended gauge symmetries with the Schwinger-Dyson symmetries through the use of collective fields. We again shift the fundamental fields ϕ_A , and we choose to include the internal gauge (BRST) transformations in the transformation of the collective fields only. The combined extended symmetries then take the form:

$$\begin{aligned}\delta_a \phi_A &= \phi_{Aa}^* \\ \delta_a \bar{\phi}_A &= \phi_{Aa}^* - \mathcal{R}_{Aa}^{(\phi-\bar{\phi})} \\ \delta_a \phi_{Ab}^* &= \epsilon_{ab} B_A \\ \delta_a B_A &= 0\end{aligned}\tag{14}$$

Gauge fixing the collective fields to zero is again done by adding a term manifestly invariant under the extended BRST symmetry,

$$\begin{aligned}\frac{1}{4}\epsilon^{ab}\delta_a\delta_b[\bar{\phi}_A(-1)^{\epsilon_B}M_{AB}\bar{\phi}_B] &= \epsilon^{ab}\phi_{Aa}^*M_{AB}\mathcal{R}_{Bb}^{(\phi-\bar{\phi})} \\ &\quad - \frac{1}{2}(-1)^{\epsilon_B}\epsilon^{ab}\bar{\phi}_A M_{AB} \frac{\delta^r \mathcal{R}_{Bb}^{(\phi-\bar{\phi})}}{\delta\phi_C} \mathcal{R}_{Ca}^{(\phi-\bar{\phi})} \\ &\quad + \bar{\phi}_A(-1)^{\epsilon_B+1}M_{AB}B_B - \frac{1}{2}\epsilon^{ab}\phi_{Aa}^*M_{AB}\phi_{Bb}^* \\ &\quad - \frac{1}{2}\epsilon^{ab}\mathcal{R}_{Aa}^{(\phi-\bar{\phi})}M_{AB}\mathcal{R}_{Bb}^{(\phi-\bar{\phi})}\end{aligned}\tag{15}$$

Notice that the “antifields” ϕ_{Aa}^* (really the ghost-antighost pair of the Schwinger-Dyson shift symmetries) automatically start acting as sources for the internal BRST–anti-BRST symmetries, while the collective fields $\bar{\phi}$ acts as a source for the commutator of the two.

The extended action also contains terms quadratic in the ghost-antighost fields ϕ_{Aa}^* . Of course, when we integrate over these fields (as well as over $\bar{\phi}_A$ and B_A), we simply recover the starting action S integrated over the standard measure. The formulation in these terms is therefore entirely consistent, and we can now finally add a gauge-fixing term of the form $\epsilon^{ab}\delta_a\delta_b\chi(\phi)$. The interesting effect of adding such a term is that the collective field $\bar{\phi}$ is no longer set equal to zero in the gauge-fixing procedure. The resulting gauge-fixed action seems at first sight to imply very unusual Feynman rules since, *e.g.*, the argument of even the classical action is shifted by $\delta^r\chi(\phi)/\delta\phi_A$. It can, however, formally be brought into the conventional form [6] by a field redefinition. This aspect is avoided entirely if we choose the “gauge-boson” χ to be a function of the difference $\phi_A - \bar{\phi}_A$ instead. The gauge-fixing is then directly identical to the standard procedure. On the other hand, no new insight into the gauge-fixing has been gained.

Instead, it is convenient to separate out the terms of the extended action that act as sources for the BRST–anti-BRST transformations of the internal symmetries, and their commutator. We therefore define one part S_{BLT} of the extended action S_{ext} through

$$S_{ext} = S_0(\phi - \bar{\phi}) + \frac{1}{4}\epsilon^{ab}\delta_a\delta_b[\bar{\phi}_A(-1)^{\epsilon_B}M_{AB}\bar{\phi}_B]$$

$$\begin{aligned}
&= S_{BLT} + \bar{\phi}_A (-1)^{\epsilon_B+1} M_{AB} B_B - \frac{1}{2} \epsilon^{ab} \phi_{Aa}^* M_{AB} \phi_{Bb}^* - \frac{1}{2} \epsilon^{ab} \mathcal{R}_{Aa} M_{AB} \mathcal{R}_{Bb} \\
&= S_{BLT} + S_{aux}
\end{aligned} \tag{16}$$

Clearly, S_{BLT} is nothing but the classical action plus the source terms for the internal extended BRST transformations. We have called what is left (the quadratic part of the ghost-antighost fields ϕ_{Aa}^* , the term that fixes the collective field to zero, and the term independent of ϕ_{Aa}^* and $\bar{\phi}_A$) the “auxiliary action”, and denoted it by S_{aux} . It is auxiliary in the sense that in the end it only serves to remove the $\phi_{Aa}^*, \bar{\phi}_A$ -fields from the path integral in a correct manner. The reason for this splitting-up of the extended action is twofold. First, it allows us to formulate the construction of S_{ext} by means of a differential equation reminiscent of the conventional Master Equations [4, 3]. Second, it allows us to perform the subsequent fixing of the original gauge symmetries in an elegant way, uncovering a remnant of the gauge-fixing procedure of Batalin and Vilkovisky [4]. At this final step we will make complete contact to the previous work of Batalin, Lavrov and Tyutin [7].

Let us first discuss the Master Equations. At the level where the collective fields and the auxiliary fields B_A are still kept in the action, these Master Equations take rather trivial forms. They simply express invariance of the extended action S_{ext} under both symmetries δ_a . Thus, $0 = \delta_a S_{ext}$ immediately implies

$$\frac{\delta^r S_{ext}}{\delta \phi_A} \phi_{Aa}^* + \frac{\delta^r S_{ext}}{\delta \bar{\phi}_A} (\phi_{Aa}^* - \mathcal{R}_{A+}^{(\phi-\bar{\phi})}) + \frac{\delta^\nabla \mathcal{S}_{|\S\sqcup}}{\delta \phi_{A\downarrow}^*} \varepsilon_{\downarrow\downarrow} \mathcal{B}_A = 0. \tag{17}$$

But inserting the decomposition (16) into these equations give us more conventional-looking Master Equations for the source part S_{BLT} . (The easiest way to derive it is to note that we are now, after gauge-fixing $\bar{\phi}_A$ to zero, free to shift the fields $\phi_A \rightarrow \phi_A + \bar{\phi}_A$.) One finds

$$\frac{\delta^r S_{BLT}}{\delta \phi_A} \mathcal{R}_{Aa} + \frac{\delta^r S_{BLT}}{\delta \bar{\phi}_A} \phi_{Aa}^* = 0. \tag{18}$$

Then, using the fact that S_{BLT} contains the sources for the internal BRST–anti-BRST transformations, we note that it can be rewritten as

$$\frac{\delta^r S_{BLT}}{\delta \phi_A} \varepsilon_{ab} M_{AC}^{-1} \frac{\delta^l S_{BLT}}{\delta \phi_{Cb}^*} + \frac{\delta^r S_{BLT}}{\delta \bar{\phi}_A} \phi_{Aa}^* = 0. \tag{19}$$

After a linear field redefinition,

$$\phi_{Aa}^* = \varepsilon_{ba} \phi_B'^{*b} M_{BA}^{-1}, \quad \bar{\phi}_A = (-1)^{\epsilon_B} \bar{\phi}_B' M_{BA}^{-1}, \tag{20}$$

this yields

$$\frac{\delta^r S_{BLT}}{\delta \phi_A} \frac{\delta^l S_{BLT}}{\delta \phi_A'^{*a}} = \varepsilon_{ab} \frac{\delta^l S_{BLT}}{\delta \bar{\phi}_A'} \phi_A'^{*b}. \tag{21}$$

We have for convenience dropped primed indices on the new variables. The l.h.s. of eq. (21) is nothing but half the generalized antibracket $(S, S)_a$. Notice that after the

field-relabelling the matrix M has disappeared from S_{BLT} , which now takes the simple form

$$S_{BLT} = S_0(\phi) + \phi_A^{*a} \mathcal{R}_{A+} + \frac{\infty}{\in} \frac{\delta^\nabla \mathcal{R}_{A+}}{\delta \phi_B} \mathcal{R}_{B|} . \quad (22)$$

Due the redefinition of what we now view as ghost-antighosts ϕ_A^{*a} and collective fields $\bar{\phi}_A$, the ghost number assignments have changed. One easily sees that now

$$gh(\phi_A^{*a}) = (-1)^a - gh(\phi^A) , \quad gh(\bar{\phi}_A) = -gh(B_A) = -gh(\phi^A) , \quad (23)$$

while of course Grassmann parities remain unchanged.

Gauge-fixing in a manifest BRST-anti-BRST invariant manner is again done by adding a term of the kind

$$\begin{aligned} \frac{1}{2} \varepsilon^{ab} \delta_a \delta_b \chi(\phi) &= \frac{1}{2} \varepsilon^{ab} \frac{\delta^r \chi(\phi)}{\delta \phi_B} \frac{\delta^r \mathcal{R}_{B|}}{\delta \bar{\phi}_A} \mathcal{R}_{A+} + \frac{\infty}{\in} \varepsilon^{-|} (-\infty)^{\varepsilon_B} \frac{\delta^\nabla}{\delta \phi_A} \frac{\delta^\nabla}{\delta \phi_B} \chi(\phi) \mathcal{R}_{A+} \mathcal{R}_{B|} \\ &= -\frac{\delta^r \chi(\phi)}{\delta \phi_A} \frac{\delta^l S_{BLT}}{\delta \bar{\phi}_A} + \frac{1}{2} \varepsilon^{ab} (-1)^{\varepsilon_B} \frac{\delta^r}{\delta \phi_A} \frac{\delta^r}{\delta \phi_B} \chi(\phi) \frac{\delta^l S_{BLT}}{\delta \phi_A^{*a}} \frac{\delta^l S_{BLT}}{\delta \phi_B^{*b}} \end{aligned} \quad (24)$$

The partition function can then be written

$$\begin{aligned} \mathcal{Z} &= \int [d\phi_A] [d\phi_A^*] [d\bar{\phi}_A] e^{\left[\frac{i}{\hbar} S_{aux} + S_{gf} + S_{BLT} \right]} \\ &= \int [d\phi_A] [d\phi_A^*] [d\bar{\phi}_A] e^{\frac{i}{\hbar} S_{aux}} \left[\hat{U} e^{\frac{i}{\hbar} S_{BLT}} \right] , \end{aligned}$$

with a “gauge-fixing operator” defined by

$$\hat{U} \equiv \exp \left[-\frac{\delta^r \chi(\phi)}{\delta \phi_A} \frac{\delta^l}{\delta \bar{\phi}_A} - \frac{i\hbar}{2} \varepsilon^{ab} \frac{\delta^l}{\delta \phi_B} \frac{\delta^r}{\delta \phi_A} \chi(\phi) \frac{\delta^l}{\delta \phi_A^{*a}} \frac{\delta^l}{\delta \phi_B^{*b}} \right] . \quad (25)$$

The Master equation (21) for S_{BLT} corresponds to the one proposed by Batalin, Lavrov and Tyutin [7], and the gauge-fixing procedure through the matrix $\hat{U}$ is also identical to theirs⁷. The only difference is that from the present derivation the ghost-antighost pair ϕ_{Aa}^* are not set equal to zero after having operated with $\hat{U}$ on S_{BLT} . Rather, these fields are now integrated over with the “measure”

$$d\mu[\phi^*] \equiv [d\phi^*] e^{\left[\frac{i}{\hbar} S_{aux} \right]} , \quad (26)$$

but it is easy to see that the end result is the same. The collective fields $\bar{\phi}_A$ are precisely set equal to zero after the integration over B_A in S_{aux} . Through this derivation, we have thus managed to stipulate in a very precise manner what is behind this gauge-fixing procedure based on $\hat{U}$, and how the fields $\phi_{Aa}^*, \bar{\phi}_A$ are to be treated in the path integral. It is remarkable that the whole formalism originally could be arrived at through algebraic considerations alone [7], apparently only guided by the principle of treating the “antifields” as sources for the internal BRST and anti-BRST symmetries (and their

⁷In the form given by Lavrov [10].

independent combination in their commutator), and some analogies with the Hamiltonian formalism [11].

We see now that the mysterious three sets of “antifields” ϕ_{Aa}^* and $\bar{\phi}_A$ are nothing but linear combinations of the ghost-antighost pairs of ref. [3], and the collective fields used to derive the Schwinger-Dyson BRST symmetry, respectively. We now also understand the reason why these collective fields can not immediately be disposed of, as in the case of keeping just BRST (or anti-BRST) symmetry: The lack of nilpotency of the operator $\varepsilon^{ab}\delta_a\delta_b$ after having integrated out the collective fields and their auxiliary fields (B_A). If one integrates out *only* the collective fields, while keeping the B_A -fields, nilpotency of the extended BRST operators may be kept. But the resulting formalism does not appear very convenient.

As in the case of just BRST or anti-BRST symmetries, one sees that the Master Equation must be satisfied due to much more general principles than independence of the gauge-fixing functions. Also the Master Equations of *extended* BRST symmetries ensure that correct Schwinger-Dyson equations are obtained even after the subsequent gauge-fixing of internal symmetries.

The fact that the partition function is independent of the operator $\hat{U}$ follows in an entirely straightforward manner from the present perspective: It simply reflects the addition of an extended-BRST-exact term to the action. Since during the introduction of this operator we have kept all fields, the operator $\varepsilon^{ab}\delta_a\delta_b$ is nilpotent when acting on an arbitrary function of all the fields $\phi_A, \phi_{Aa}^*, \bar{\phi}_A$. The form of $\hat{U}$ derived here is thus very specific for gauge bosons χ that only depend on ϕ_A . The general expression for $\varepsilon^{ab}\delta_a\delta_b$ is straightforward to write out. Such a term can be added to the action without changing the physical content, but the resulting actions can look highly unusual. The addition of such a general term $\varepsilon^{ab}\delta_a\delta_b$ to the action (and the analogous modified $\hat{U}$ -operation) has a direct counterpart in the general operator

$$\hat{U} = e^{i\hbar\hat{T}(X)} , \quad \hat{T} = \frac{1}{2}\varepsilon_{ab}[\bar{\Delta}^b, [\bar{\Delta}^a, X]_-]_+ \quad (27)$$

which is discussed in refs. [7, 10]. Here $\bar{\Delta}^a \equiv \Delta^a + (i/\hbar)\varepsilon^{ab}\phi_{Ab}^*\delta^l/\delta\bar{\phi}_A$. The operator $\hat{T}$ is thus apparently the closest one can get to the analogue of the “quantum BRST-anti-BRST operator” equivalent of $\varepsilon^{ab}\delta_a\delta_b$.

It should be very clear from our presentation that – as in the case of just BRST or anti-BRST symmetries – there is no unique Lagrangian quantization prescription. In the end we formulated the collective field quantization in a manner that directly proved its equivalence to the scheme of Batalin, Lavrov and Tyutin [7], but there are obviously many ways of departing from that particular formulation.

Since the Master Equation (21) has been *derived* from known principles using the collective field technique, its present use may seem rather limited. What is needed in order to go beyond the derivation in terms of collective fields is an overall formulation that ensures correct Schwinger-Dyson equations as Ward Identities, without losing nilpotency of the operator $\varepsilon^{ab}\delta_a\delta_b$. As long as the derivation is directly tied to the collective fields, the answer in a sense needs to be known beforehand. One can nevertheless formulate empirical rules and boundary conditions for the Master Equations (21) that will guar-

antee the correct prescription for the gauge-fixing. Whether or not theories with open gauge algebras can be made invariant under extended BRST symmetries (a problem that from the present point of view requires that we go beyond the relatively straightforward collective field technique used here), and how it ties up with Master Equations related to those of eq. (21) remains to be investigated.

Related to the close connection with collective fields is the question of quantum corrections to the Master Equations. Such terms of the kind $\Delta^a S$ do not appear when the derivation can be phrased in terms of collective fields in the manner described here, because then the functional measure is (formally) guaranteed to be invariant. Ultraviolet regulators that break some of the symmetries may correspond to non-invariant measures, and then these terms should be included.

Although the Master Equations (21) can be written in terms of the generalized antibracket, it is clear that the canonical structure present when quantizing with either BRST or anti-BRST symmetry alone is lost. This is entirely natural from the point of view of ref. [3], since the full canonical structure only appears when one integrates out one of the fields ϕ_{Aa}^* , while keeping its ghost or antighost partner in the path integral. In this manifestly BRST–anti-BRST invariant formulation it cannot be achieved, and the canonical formalism is therefore not fully present. The same holds for the “quantum BRST” (or anti-BRST) operator, which is simply the Schwinger-Dyson BRST (anti-BRST) operator after the integration of only one of the ghost-antighosts ϕ_{Aa}^* [3]. Such forms of the operators have no natural rôles in our $Sp(2)$ invariant formulation either, and the operator $\hat{T}$ above is probably the closest one can get. It is, however, not a priori ruled out that the extended BRST symmetries can be introduced only based on the Schwinger-Dyson BRST operator. Although $Sp(2)$ symmetry will not be manifest at all levels, it may perhaps be imposed on the subspace of fields ϕ^A , which, after all, would be sufficient.

The geometric interpretation of the usual Batalin-Vilkovisky quantization scheme has recently received considerable attention [12]. In an interesting paper, Henneaux [13] has shown that several aspects of this geometric interpretation can be carried over to the case of extended BRST symmetries when based on the formulation given in ref. [7]. An obvious open problem is to give a geometric interpretation of the new ghosts c^A of ref. [3], without being restricted to an $Sp(2)$ -invariant formulation.

References

- [1] J. Alfaro and P.H. Damgaard, Phys. Lett. **B222** (1989) 425.
J. Alfaro, P.H. Damgaard, J. Latorre and D. Montano, Phys. Lett. **B233** (1989) 153.
- [2] J. Alfaro and P.H. Damgaard, Ann. Phys.(N.Y.) **202** (1990) 398.
- [3] J. Alfaro and P.H. Damgaard, preprint CERN-TH-6788/93, hep-th 9301103.
- [4] I.A. Batalin and G.A. Vilkovisky, Phys. Lett. **B102** (1981) 27; Phys. Rev. **D28** (1983) 2567 [E: **D30** (1984) 508]; Nucl. Phys. **B234** (1984) 106; J. Math. Phys. **26** (1985) 172.
- [5] M. Henneaux, Nucl. Phys. B (Proc. Suppl.) **18A** (1990) 47.
J.M.L. Fisch, Univ. Brux. preprint ULB TH2/90-01.
W. Troost and A. Van Proeyen, lecture notes, Leuven Univ. Press. (to appear).
- [6] G. Curci and R. Ferrari, Phys. Lett. **B63** (1976) 91.
I. Ojima, Prog. Theor. Phys. Suppl. **66** (1979) 324.
N. Nakanishi and I. Ojima, Z. Phys. **C6** (1980) 155.
L. Alvarez-Gaumé and L. Baulieu, Nucl. Phys. **B212** (1983) 255.
F.R. Ore and P. van Nieuwenhuizen, Nucl. Phys. **B204** (1982) 317.
- [7] I.A. Batalin, P.M. Lavrov and I.V. Tyutin, J. Math. Phys. **31** (1990) 1487; **32** (1991) 532; **32** (1991) 2513.
- [8] J. Gomis and J. Roca, Nucl. Phys. **B343** (1990) 152.
C.M. Hull, Mod. Phys. Lett. **A5** (1990) 1871.
- [9] W. Troost, P. van Nieuwenhuizen and A. Van Proeyen, Nucl. Phys. **B333** (1990) 727.
F. De Jonghe et al., Phys. Lett. **B289** (1992) 354.
- [10] P.M. Lavrov, Mod. Phys. Lett. **A6** (1991) 2051.
- [11] I.A. Batalin, P.M. Lavrov and I.V. Tyutin, J. Math. Phys. **31** (1990) 6.
I.A. Batalin and E.S. Fradkin, Ann. Inst. H. Poincaré (Phys. Theor.) **49** (1988) 145.
- [12] E. Witten, Mod. Phys. Lett. **A5** (1990) 487.
A. Schwarz, UC Davis preprint (1992), hep-th 9205088.
E. Witten, preprint IASSNS-HEP-92/63, hep-th 9208027.
H. Hata and B. Zwiebach, preprint MIT-CTP-2177, hep-th 9301097.
- [13] M. Henneaux, Phys. Lett. **B282** (1992) 372.